1 Long-distance benefits of marine reserves: myth or reality?

2 Stéphanie Manel¹, Nicolas Loiseau², Marco Andrello³, Katharina Fietz⁴, Raquel Goñi⁵, Aitor
3 Forcada⁶, Philippe Lenfant⁷, Stuart Kininmonth^{8,9}, Concepción Marcos¹⁰, Virginie Marques²,
4 Sandra Mallol⁵, Angel Pérez-Rufaza¹⁰, Corinna Breusing⁴, Oscar Puebla⁴, David Mouillot^{2,11}

5

6 1-EPHE, PSL Research University, CEFE UMR 5175, CNRS, Université de Montpellier,
7 Université Paul-Valéry Montpellier, Biogéographie et Ecologie des Vertébrés, 1919 route de
8 Mende, Montpellier 34293, France

9 2-UMR 9190 MARBEC, IRD-CNRS-IFREMER-UM, Université de Montpellier, Montpellier
10 34095 France

11 3- Department of Ecology and Evolutionary Biology, University of Toronto, Toronto,
12 Ontario, Canada

13 4- GEOMAR Helmholtz Centre for Ocean Research Kiel, Evolutionary Ecology of Marine
14 Fishes, Düsternbrooker Weg 20, 24105 Kiel, Germany.

15 5- Instituto Español de Oceanografía, Centro Oceanográfico de Baleares, Moll de Ponent s/n,
16 07015 Palma de Mallorca, Spain

17 6- Department of Marine Sciences and Applied Biology, University of Alicante, P.O. Box 99,
18 03080 - Alicante, Spain

19 7- Univ. Perpignan Via Domitia, Centre de Formation et de Recherche sur les
20 Environnements Méditerranéens, UMR 5110, 58 Avenue Paul Alduy, F-66860 Perpignan,
21 France

22 8- School of Marine Studies, University of South Pacific, Suva, Fiji

23 9- Stockholm Resilience Centre, Stockholm University, Stockholm, Sweden

24 10- Departamento de Ecología e Hidrología, Facultad de Biología, Campus de Espinardo,
25 Regional Campus of International Excellence “Campus Mare Nostrum”, University of
26 Murcia, 30100 Murcia, Spain

27 11-Australian Research Council Centre of Excellence for Coral Reef Studies, James Cook
28 University, Townsville, QLD 4811 Australia.

29 **Correspondence: stephanie.manel@ephe.psl.eu (S. Manel)**

30

31 **Keywords:** connectivity, global network, long-distance dispersal, marine reserves, marine
32 protected areas.

33
34 **Abstract**

35
36 Long-distance (>40km) dispersal from marine reserves is poorly documented. Yet, it can
37 provide essential benefits such as seedling fished areas or connecting marine reserves into
38 networks. From a meta-analysis, we suggest that the spatial scale of marine connectivity is
39 underestimated due to the limited geographic extent of sampling designs. We also found that
40 the largest marine reserves (>1,000km²) are the most isolated. These findings have important
41 implications for the assessment of evolutionary, ecological and socio-economic long-distance
42 benefits of marine reserves. We conclude that existing methods to infer dispersal should
43 consider the up-to-date genomic advances and also expand the spatial scale of sampling
44 designs. Incorporating long-distance connectivity in conservation planning will contribute to
45 increase the benefits of marine reserve networks.

56

57

58

59 **Benefits from marine reserves: where are we?**

60 Marine resources are declining at an alarming rate [1, 2], with more than half of the oceanic
61 area exploited by industrial fishing [3]. In response, marine protected areas (MPAs) have been
62 established in an effort to conserve biodiversity and sustain fisheries [4-7]. Yet, only 3.7 % of
63 the ocean is presently covered by MPAs and less than 2% by no-take MPAs (referred to as
64 marine reserves [8]) specifically (mpatlas.org; [9]). Despite the recent establishment of large-
65 scale MPAs ($>100,000 \text{ km}^2$) [10], the current trend of protection is not keeping pace with the
66 increasing human footprint on marine resources [11, 12]. In particular, the Aichi Biodiversity
67 Target 11 established by the Convention of Biological Diversity to protect at least 10% of the
68 ocean by 2020 is unlikely to be achieved [9, 13, 14]. In consideration of the ever-growing
69 human population, fishing technological developments and per capita consumption rates, a
70 new target of 30% protected area by 2030 was proposed at the 2016 International Union for
71 Conservation Nature (IUCN) World Conservation Congress in line with scientific advice [6,
72 15]. There is thus an urgent need to better understand the full range of benefits provided by
73 marine reserves to optimize future conservation efforts.

74

75 Theoretical and empirical studies support the positive effects of marine reserves within their
76 boundaries and in their vicinity [16-19]. Indeed, marine reserves unambiguously host more
77 and larger - and thus more fertile - individuals than fished areas (e.g. [5, 20, 21]). They also
78 contribute to preserve genetic diversity [22], increase human wellbeing [23], alleviate poverty
79 [24] and facilitate adaptation to climate change [25]. In addition, when properly designed and
80 enforced, they have the potential to increase catches of commercial species in surrounding

81 fishing grounds [26, 27] due to juvenile or adult **spillover** (see glossary) [28, 29]. However,
82 such direct benefits have been typically documented at short distances from reserve
83 boundaries, i.e. from a few hundred meters to less than 40 km [16, 30 , 31]. In contrast, little
84 is known about the benefits of marine reserves in areas that are 40 to hundreds of kilometers
85 away from their boundaries [32].

86

87 The concept of long-distance (>40 km) **dispersal** is not new in marine ecology and the oceans
88 have been assumed to function as mostly open, well-connected systems until the last two
89 decades [33, 34], when evidence of local recruitment started to accumulate [35, 36]. However,
90 the large-scale impacts of marine reserves resulting from long-distance dispersal of larvae
91 [37], juveniles and adults [38] are still poorly documented. The potential for long-distance
92 dispersal is highest for **pelagic species**, which is consistent with their widespread geographic
93 distributions [39]. Yet, recent findings based on telemetry and genetic tools indicate that
94 **benthic** and **demersal species** (hereafter called benthos), including those in the deep sea, can
95 also disperse up to hundreds of kilometers [32, 40-42]. This recognition of high dispersal
96 capabilities calls for more studies on the effect of dispersal far from reserves and for a better
97 integration of long-distance dispersal in the design of reserve networks.

98

99 Here we review the potential long-distance benefits of marine reserves, including those
100 provided by relatively rare long-distance dispersal events. We focus on the benthos since their
101 adult stages can be more easily assigned to protected versus non-protected areas compared to
102 pelagic species which have large home ranges, often larger than most MPAs [43]. First,
103 through an extensive literature screening, we characterize the spatial scale of dispersal for the
104 benthos. We then discuss how existing methods can be enhanced to expand the scale of
105 connectivity analyses. Finally, we discuss the potential long-distance benefits of marine

reserves for both conservation and fisheries and how a well-connected network can enhance those benefits.

Spatial scale of dispersal and connectivity in the marine realm: do we capture the full picture?

We define marine connectivity as the exchange of individuals among marine populations [44]. This exchange can take place through dispersal of individuals as larvae, juveniles, or adults. When individuals reproduce successfully thereafter, **demographic connectivity** translates into **genetic connectivity**. Determining the spatial scale of marine connectivity is crucial for our understanding of the population dynamics, genetic structure and biogeography of marine organisms, and accordingly for the design of marine reserves.

To obtain a global estimate of the spatial scale of marine connectivity for the benthos, we conducted an extensive – but non-exhaustive – literature review over the last decade in the ISI Web of Science (supplementary text S1, supplementary file S1). Of the 460 papers identified, 130 were included in our meta-analysis as they contain information about maximum sampling geographic range and maximum inferred demographic or genetic connectivity for a total of 243 species.

The different methods used to estimate dispersal distance apply to different spatial and temporal scales (Table I of Box 1). The median potential dispersal distance averaged across all studies based on biophysical models (226 km, interquartile range = 160 - 415 km, number of species = 56) was at least four times higher than the demographic (realized and effective) median dispersal distance (42 km, interquartile range = 27 – 250 km, number of species = 55;

Figure 1A; Box 1). These results indicate that studies are either overestimating potential dispersal or underestimating demographic dispersal, with some exceptions where both estimates are congruent (e.g. [45]). However, assessing the full spatial extent of dispersal is challenging due to the inherent difficulty of tracking or recapturing organisms over long distances. With a few exceptions (e.g. [32, 41, 46]), most empirical studies of demographic connectivity were conducted at scales smaller than 40 km [47]. Estimating demographic connectivity at larger spatial scales and over multiple generations would require sampling significantly more individuals and in more distant populations, which would entail high, possibly prohibitive, costs. Genetic assignment approaches at the population level might be scaled-up more easily than mark-recapture or parentage analysis methods and constitute a promising approach when populations are genetically differentiated [48, 49]. For example, putative first-generation migrants between two populations separated by 400 km were detected in the Omani Clownfish (*Amphiprion omanensis*) using assignment tests [40]. If populations are locally adapted, the use of genetic markers that are under divergent selection can contribute to increase the power of such approaches, and can even be used in the absence of neutral genetic structure [50]. Genetic **isolation by distance** at the population or individual level [51] provides dispersal estimates that are consistent with demographic dispersal estimates obtained from parentage analysis (Box 1) [52]. When a reference genome and **haplotype** data are available, the consideration of **admixture tracts** [53] and **blocks of identity by descent** [54] constitutes another promising avenue to detect recent dispersal events, that can also apply in isolation by distance contexts [55].

Our literature review also reveals that genetic connectivity, based on Wright's Fixation Index (F_{ST}), tends to provide higher estimates than any other method (Figure 1A: median = 910 km, interquartile range = 315 - 2346 km, number of species = 126). Yet, genetic connectivity

differs from demographic connectivity as it integrates the effects, not just off migration, but also genetic drift, mutation and selection. Translating genetic connectivity into demographic estimates of dispersal is not straightforward [56]. This notably implies estimating effective population sizes [57] or assuming specific population genetic models that are often unrealistic in real-world situations [58]. Furthermore, **gene flow** over large geographic distances might result from **stepping-stone dispersal** over multiple generations without necessarily implying direct long-distance dispersal events [59].

Globally, the data show a universal positive correlation between the geographic sampling scale of the study and the maximum dispersal or connectivity averaged across all studies and organisms ($R = 0.7$, $p < 0.001$; Figure 1B). The relation holds true when analyzing the data per type of dispersal estimate (potential vs. demographic vs. genetic). In 45 % of the studies, the dispersal distance was equal to the maximum geographic extent of the sampling. This reached 48 % when data were restricted to coastal fishes, 41 % for invertebrates and 41 % for deep sea organisms (Figure 1C). These results suggest that estimates are limited by the spatial scale of the sampling, resulting in a global underestimation of the extent of demographic and genetic connectivity.

For genetic connectivity, an absence of population structure can also result from a lack of statistical power to detect subtle population genetic structure when a small number of genetic markers are used. With the advent of **next-generation sequencing** technologies, this limitation can now be overcome by typing hundreds to millions of **single nucleotide polymorphism markers** [60, 61].

A variety of mechanisms can contribute to long-distance dispersal in marine ecosystems (Figure 2). Foremost, the hydrodynamic forces at play in the marine environment are expected to have a strong influence on the dispersal of **pelagic larvae** [62]. In addition, the conditions encountered in the pelagic environment might influence growth, survival and pelagic larval duration, all of which can in turn induce extreme values in spatial and temporal connectivity patterns. Active larval behavior can also play an important role for the benthos [63, 64]. Extreme events, such as tsunamis [65], and oceanographic eddies and fronts [66] are also important, but overlooked potential dispersal vectors over long distances. They can favor the survival and establishment of individuals beyond their usual dispersal range. Furthermore, marine debris of natural or anthropogenic origin can constitute effective oceanic rafts for dispersal [67, 68]. These debris provide refuges for larvae and adults of **sessile species**, allowing the movement and potential establishment of a variety of species over large distances. For instance, mussels from Japan arrived on the west coast of the US after nearly six years at sea on debris produced by the 2011 East Japan earthquake [65]. Ice blocks also allow invertebrates to disperse across distances of about 20 km per day [69]. Anthropogenic vectors such as international vessel traffic, restocking from aquaculture [70], and species translocation (Box 2) also have the potential to modify the natural spatial and temporal patterns of marine connectivity [71]. Overall, the occurrence of long-distance dispersal events, even if rare, suggest that marine reserves can have an effect far beyond their boundaries, which calls for a re-evaluation of the spatial extent of their potential benefits.

Long-distance benefits of marine reserves

We consider a long-distance benefit of marine reserve any change in biomass, biological processes (e.g. recruitment) or biodiversity (including genetic diversity as raw material for adaptation) at distance greater than 40 km from reserve boundaries that contributes to

improve ecosystem function or human livelihoods (e.g. fisheries, tourism, culture) (Figure 2). Long-distance benefits from reserves can occur at different spatial and temporal scales depending on the vector of dispersal (Box 1, Figure 2), and this includes both direct and stepping-stone dispersal processes.

Parentage analyses have demonstrated dispersal from marine reserves at more than 40 km [32, 46]. For example, Almany *et al.* [46] revealed connectivity patterns with direct exchanges of larvae over up to 150 km among reefs with varying levels of protection. **Larval dispersal** from reserves towards exploited areas located at more than 100 km has also been suggested by biophysical models [37, 72]. Yet, empirical studies showing an effect of marine reserves on fished areas are largely restricted to spatial scales smaller than 40 km (e.g. [26, 73, 74]). Scaling-up these studies is challenging for a variety of logistic reasons, including the difficulty to sample and monitor individuals over large spatial scales.

Even if long-distance dispersal events from marine reserves are rare, a few successful migrants can be sufficient to re-colonize areas where local populations have been extirpated or to expand species distributions in response to global change [75]. Long-distance dispersal between populations that are genetically differentiated or locally adapted can moreover contribute to limit inbreeding, increase genetic diversity and facilitate adaptation to a changing environment [25, 76]. However, in case of local adaptation, long-distance dispersal can also reduce fitness of recipient populations through immigration of locally maladapted alleles [77].

Active translocations from marine reserves can also contribute to restore locally depleted or extinct populations [78]. For example, the Bumphead Parrotfish (*Bolbometopon muricatum*)

is highly targeted by spear fishers due to its large size and therefore population densities tend to be low in areas close to human populations [79]. This species is only abundant in reserves and on the most protected reefs such as in Palau where it aggregates to spawn. These populations have been used as a source of eggs and larvae for active translocations (Box 2). Nearly 500 translocation projects of 242 marine species have been recorded [78]. However, still few projects take advantage of large stocks in marine reserves. It should also be emphasized that translocations entail a number of risks (e.g. disease, invasion, gene pool mixing). Overall, marine reserves could support a wide variety of long-distance benefits that are potentially underestimated and that should be considered for the design of reserve networks.

Implications of long-distance dispersal for marine reserve design

An underestimate of dispersal ability can profoundly influence the design of marine reserve networks. Long-distance dispersal can potentially connect distant and isolated reserves, and sustain biodiversity and biomass in exploited areas located at more than 40 km from their boundaries. In this respect, long-distance dispersal provides a fresh perspective on two long-lasting and active debates in the marine reserve literature.

First, long-distance dispersal has implications for the unresolved single-large-or-several-small (SLOSS) marine reserve debate. Simulations suggest that a network of well-connected reserves on a scale of 10-100 km can meet both conservation and fisheries goals [7, 80]. However, the idea that a network constituted of many small reserves spaced within species maximum dispersal distance [4] maximizes reserve benefits to fisheries has been recently revisited and challenged. Based on a spatially explicit model of population dynamics, De Leo & Micheli [81] show that for larval dispersal >10 km, one or two large reserves are more

efficient in terms of fisheries gains than 10 or 20 small reserves covering the same area. This is notably due to the fact that large ($>100 \text{ km}^2$), old (>10 years, [5]) and well-managed reserves tend to increase fish density and biomass [5, 82], and that large females over-contribute to reproduction since the relation between female body mass and reproductive output is hyperallometric for the vast majority of fishes (i.e. a 2-kg female has a higher reproductive output in terms of egg number, volume and energy, than two 1-kg females [83]). We can therefore expect large, old and well-managed reserves to disproportionately contribute to larval seedling within a network when dispersal distances are greater than ten kilometers. Using a rigorously calibrated metapopulation model with empirical data from the Great Barrier Reef, Hopf et al. [84] also show that reserves are unable to compensate for the increased mortality outside reserve boundaries when they are small or at the periphery of the metapopulation [84]. In contrast, the establishment of a single large reserve, that is able to seed overexploited areas through dispersal, is expected to result in higher population growth within reserve boundaries and shorter recovery times after overexploitation outside the reserve. Finally, a global analysis indicates that when larval dispersal distances are long ($>40 \text{ km}$), the magnitude of biomass increase within large reserves is expected to be sufficient to compensate for the redistributed fishing pressure associated with reserve establishment [85].

The median nearest-neighbor distance between marine reserves is estimated at 12 km globally (interquartile distance: 4 to 40 km) (Supplementary text S2). This geographical pattern is highly variable, with some reserves being very isolated (e.g. the Parque Natural Obô do Príncipe in São Tomé and Príncipe at 4130 km from the nearest reserve, the Monumento Natural do Arquipélago de São Pedro e São Paulo at 940 km off the coast of Brazil) (Figure 3A, supplementary text S2). Fortunately, 76 % of reserves are found closer to the nearest reserve than the median demographic dispersal distance estimated in our literature review (42

km) (Figure 3B). It implies that three-quarters of marine reserves are potentially demographically embedded in a connected network providing spatial insurance. The median nearest-neighbor distance in the 24 % remaining reserves is estimated to be 129 km, with a very skewed distribution (Figure 3B). Notably, 83% of large reserves ($> 1000 \text{ km}^2$) are isolated ($>42\text{km}$) (Figure 3C). The mean nearest neighbor of this subset of large marine reserves is 359 km away, decreasing their potential contribution to the global network. Therefore, the largest marine reserves, allowing large spillover of individuals and providing benefits for both biodiversity and human population, are the least connected. However, we did not consider how sea surface currents could modify our assessment of connectivity among isolated reserves.

Second, long-distance dispersal has also implications for prioritizing the conservation of human-impacted versus non-impacted areas. Intuitively, one can see little benefit in placing reserves in isolated areas which are difficult to access and therefore *de facto* protected [86]. On the other hand, reserves close to dense human populations can mitigate but not eliminate the high anthropogenic pressure outside but also inside their boundaries [87]. It has therefore been suggested that reserves located at an intermediate level of human pressure might offer the maximum benefits in terms of fish biomass within their boundaries [87]. For top predators like sharks, only isolated marine reserves with low human pressure can be effective [87]. The realization of long-distance dispersal would also make a case for the protection of such areas isolated from human pressure. More generally, it would suggest to reconsider the design of marine reserve networks with fewer but larger reserves, including isolated ones, to sustain large populations of large individuals, even of top predators, that can massively seed larvae towards fishing grounds. Tools that integrate species dispersal in conservation planning are now available to reach both conservation and fisheries management objectives in a

multispecies framework [88]. The consideration of long-distance dispersal would certainly modify the outputs of conservation plans.

Concluding remarks

Marine dispersal has been extensively documented at short distance (mostly <40 km). We suggest that this has been due, at least in part, to logistic constraints and a restricted geographic extend of the sampling design (e.g. the median sampling distance in parentage analyses is only 33 km, interquartile range = 29 – 60 km, number of species = 22). While a significant fraction of dispersal indeed occurs at small spatial scales, the fraction of the **dispersal kernel** that we are missing is largely unknown (see Outstanding Questions). A few recent empirical studies have demonstrated dispersal of fish at larger spatial scales (up to 400 km), but even these estimates were limited by the maximum sampling distance [40]. Dispersal estimates from biophysical modelling studies at larger spatial scales suggest even longer dispersal distances (median sampling distances = 600 km, interquartile range: 237-1400, Figure 1A). Such long-distance connectivity patterns remain challenging to validate empirically, but have potentially important consequences in terms of reserve design and benefits. The more isolated reserves are, the more critical long-distance dispersal becomes to maintain source-sink dynamics between protected and exploited populations. Thus, long-distance benefits imply a more regional and network-based perspective, which entails specific challenges. Long-distance dispersal will often cross countries as well in-shore-offshore boundaries. The designation of marines reserves is largely carried out by individual countries and they rarely able coordinate efforts with other countries and high-seas authorities [89]. Furthermore, the focus on marine reserves is strongly constrained by the perspective of local stakeholders [15], which is perfectly justified but should nonetheless not obliterate a broader perspective.

We suggest scaling-up dispersal studies at regional instead of local scale. We can now genotype a large number of genetic markers, which provides the opportunity to apply population-level assignment tests at large spatial scales and in a context of low spatial structure [90]. Such studies can be guided by high-resolution biophysical models to target the specific populations among which long-distance dispersal occurs. A large number of single nucleotide polymorphism markers will also provide the statistical power to detect very subtle population structure, which will allow refining genetic connectivity estimates [90, 91]. Finally, approaches based on admixture tracts [53] and blocks of identity by descent [54] are largely untapped. The combination of genetic, chemical and biophysical approaches within an integrative statistical framework also appears to be a promising approach to estimate long-distance dispersal [92] and guide the design of new reserve networks to keep pace with ever increasing threats on marine ecosystems.

Acknowledgments

Funding and support come from the European project RESERVEBENFIT (European call BIODIVERSA3 2015-2016 call) and the CNRS for the PICS SEACONNECT. We thank Pierre Lopez for the illustration 2 of this paper. We thank Tom Bowling (Biota Aquaculture, Palau) for the pictures used in the figure in Box 2. We thank David Diaz for helpful discussions. We acknowledge the editor and the three anonymous reviewers for their comments that helped clarifying several aspects of the study.

References

- 352 1. McCauley, D.J. et al. (2015) Marine defaunation: Animal loss in the global ocean. *Science*
353 DOI: 10.1126/science.1255641.
- 354 2. Worm, B. (2016) Averting a global fisheries disaster. *Proc. Natl. Acad. Sci. U.S.A.*
355 113, 4895-4897.
- 356 3. Kroodsma, D.A. et al. (2018) Tracking the global footprint of fisheries. *Science* 359, 904-
357 907.
- 358 4. Gaines, S.D. et al. (2010) Designing marine reserve networks for both conservation and
359 fisheries management. *Proc. Natl. Acad. Sci. U.S.A.* 107, 18286-18293.
- 360 5. Edgar, G.J. et al. (2014) Global conservation outcomes depend on marine protected areas
361 with five key features. *Nature* 506, 216-220.
- 362 6. Krueck, N.C. et al. (2017) Marine reserve targets to sustain and rebuild unregulated
363 Fisheries. *Plos Biol.* 15, 1-20.
- 364 7. Lubchenco, J. and Grorud-Colvert, K. (2015) Making waves: The science and politics of
365 ocean protection. *Science* 350, 382-383.
- 366 8. Costello, M.J. and Ballantine, B. (2015) Biodiversity conservation should focus on no-take
367 Marine Reserves. *Trends Ecol. Evol.* 30, 507-509.
- 368 9. Sala, E. et al. (2018) Assessing real progress towards effective ocean protection. *Mar.*
369 *Policy* 91, 11-13.
- 370 10. O'Leary, B.C. et al. (2018) Addressing criticisms of large-scale Marine Protected Areas.
371 *BioSci.* 68, 359-370.
- 372 11. Watson, R.A. et al. (2015) Marine foods sourced from farther as their use of global ocean
373 primary production increases. *Nat. Commun.* 6, DOI: 10.1038/ncomms8365.
- 374 12. Halpern, B.S. et al. (2015) Spatial and temporal changes in cumulative human impacts on
375 the world's ocean. *Nat. Commun.* 6, DOI: 10.1038/ncomms861.
- 376 13. Tittensor, D.P. et al. (2014) A mid-term analysis of progress toward international
377 biodiversity targets. *Science* 346, 241-244.

- 378 14. Di Marco, M. et al. (2016) Global biodiversity targets require both sufficiency and
379 efficiency. *Conserv. Lett.* 9, 395-397.
- 380 15. O'Leary, B.C. et al. (2016) Effective coverage targets for ocean protection. *Conserv. Lett.*
381 9, 398-404.
- 382 16. Gell, F.R. and Roberts, C.M. (2003) Benefits beyond boundaries: the fishery effects of
383 marine reserves. *Trends Ecol. Evol.* 18, 448-455.
- 384 17. Lester, S.E. et al. (2009) Biological effects within no-take marine reserves: a global
385 synthesis. *Mar. Ecol. Prog. Ser.* 384, 33-46.
- 386 18. Stobart, B. et al. (2009) Long-term and spillover effects of a marine protected area on an
387 exploited fish community. *Mar. Ecol. Prog. Ser.* 384, 47-60.
- 388 19. Sciberras, M. et al. (2015) Evaluating the relative conservation value of fully and partially
389 protected marine areas. *Fish Fish.* 16, 58-77.
- 390 20. Diaz, D. et al. (2016) A 25-year marine reserve as proxy for the unfished condition of an
391 exploited species. *Biol. Conserv.* 203, 97-107.
- 392 21. MacNeil, M.A. et al. (2015) Recovery potential of the world's coral reef fishes. *Nature*
393 520, 341-344.
- 394 22. Perez-Ruzafa, A. et al. (2006) Effects of fishing protection on the genetic structure of fish
395 populations. *Biol. Conserv.* 129, 244-255.
- 396 23. Coulthard, S. et al. (2017) Exploring 'islandness' and the impacts of nature conservation
397 through the lens of wellbeing. *Environ. Conserv.* 44, 298-309.
- 398 24. Gurney, G.G. et al. (2014) Poverty and protected areas: An evaluation of a marine
399 integrated conservation and development project in Indonesia. *Glob. Environ. Change* 26,
400 98-107.
- 401 25. Roberts, C.M. et al. (2017) Marine reserves can mitigate and promote adaptation to
402 climate change. *Proc. Natl. Acad. Sci. U.S.A.* 114, 6167-6175.

26. Kerwath, S.E. et al. (2013) Marine protected area improves yield without disadvantaging
fishers. *Nat. Commun.* 4, DOI: 10.1126/science.
27. Di Franco, A. et al. (2016) Five key attributes can increase marine protected areas
performance for small-scale fisheries management. *Sci. Rep.* 6, DOI: 10.1038/srep38135.
28. Abesamis, R.A. and Russ, G.R. (2005) Density-dependent spillover from a marine
reserve: Long-term evidence. *Ecol. Appl.* 15, 1798-1812.
29. Goni, R. et al. (2010) Net contribution of spillover from a marine reserve to fishery
catches. *Mar. Ecol. Prog. Ser.* 400, 233-243.
30. Russ, G.R. and Alcala, A.C. (2011) Enhanced biodiversity beyond marine reserve
boundaries: The cup spillth over. *Ecol. Appl.* 21, 241-250.
31. Goni, R. et al. (2006) Spillover of spiny lobsters *Palinurus elephas* from a marine reserve
to an adjoining fishery. *Mar. Ecol. Prog. Ser.* 308, 207-219.
32. Williamson, D.H. et al. (2016) Large-scale, multidirectional larval connectivity among
coral reef fish populations in the Great Barrier Reef Marine Park. *Mol. Ecol.* 25, 6039-6054.
33. Roberts, C.M. (1997) Connectivity and management of Caribbean coral reefs. *Science*
278, 1454-1457.
34. Mora, C. and Sale, P.F. (2002) Are populations of coral reef fish open or closed? *Trends*
Ecol. Evol. 17, 422-428.
35. Jones, G.P. et al. (1999) Self-recruitment in a coral reef fish population. *Nature* 402, 802-
804.
36. Swearer, S.E. et al. (1999) Larval retention and recruitment in an island population of a
coral-reef fish. *Nature* 402, 799-802.
37. Andrello, M. et al. (2017) Global mismatch between fishing dependency and larval supply
from marine reserves. *Nat. Commun.* 8, DOI: 10.1038/ncomms16039.

38. Christie, M.R. (2010) Parentage in natural populations: novel methods to detect parent-offspring pairs in large data sets. *Molecular Ecology* *Mol. Ecol. Resour.* 10, 115-128.
39. Costello, M.J. et al. (2017) Marine biogeographic realms and species endemism. *Nat. Commun.* 8, <https://doi.org/10.1038/s41467-017-01121-2>.
40. Simpson, S.D. et al. (2014) Long-distance dispersal via ocean currents connects omani clownfish populations throughout entire species range. *PLoS ONE* 9, DOI: 10.1371/journal.pone.0107610.
41. Christie, M.R. et al. (2010) Larval connectivity in an effective network of Marine Protected Areas. *PLoS ONE* 5, DOI: 10.1371/journal.pone.0015715.
42. Espinoza, M. et al. (2015) Contrasting movements and connectivity of reef-associated sharks using acoustic telemetry: implications for management. *Ecol. Appl.* 25, 2101-2118.
43. Daly, R. et al. (2018) Refuges and risks: Evaluating the benefits of an expanded Marine Protected Area network for mobile apex predators. *Divers. Distrib.* 24, 1217-1230.
44. Cowen, R.K. et al. (2006) Scaling of connectivity in marine populations. *Science* 311, 522-527.
45. Costello, M.J. (2009) How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *Proc. Royal. Soc. Lond. B Biol. Sci.* 276, 3385-3394.
46. Almany, G.R. et al. (2017) Larval fish dispersal in a coral-reef seascape. *Nat. Ecol. Evol.* 1, DOI: 10.1038/s41559-017-0148.
47. Hameed, S.O. et al. (2016) Inverse approach to estimating larval dispersal reveals limited population connectivity along 700 km of wave-swept open coast. *Proc. Royal. Soc. Lond. B Biol. Sci.* 283, doi: 10.1098/rspb.2016.0370.
48. Paetkau, D. et al. (1995) Microsatellite analysis of population-structure in Canadian polar bears. *Mol. Ecol.* 4, 347-354.

49. Manel, S. et al. (2005) Assignment methods: matching biological questions with appropriate techniques. *Trends Ecol. Evol.* 20, 136-142.
50. Gagnaire, P.-A. et al. (2015) Using neutral, selected, and hitchhiker loci to assess connectivity of marine populations in the genomic era. *Evol. Appl.* 8, 769-786.
51. Puebla, O. et al. (2012) On the spatial scale of dispersal in coral reef fishes. *Mol. Ecol.* 21, 5675-5688.
52. Pinsky, M. et al. (2017) Marine dispersal scales are congruent over evolutionary and ecological time. *Curr. Biol.* 27, 1-6.
53. Liang, M. and Nielsen, R. (2014) The lengths of admixture tracts. *Genet.* 197, <https://doi.org/10.1534/genetics.114.162362>.
54. Browning, S.R. and Browning, B.L. (2012) Identity by descent between distant relatives: detection and applications. *Annu. Rev. Genet.* 46, 617-633.
55. Ringbauer, H. et al. (2017) Inferring recent demography from isolation by distance of long shared sequence blocks. *Genet.* 205, <https://doi.org/10.1534/genetics.116.196220>.
56. Lowe, W.H. and Allendorf, F.W. (2010) What can genetics tell us about population connectivity? *Mol. Ecol.* 19, 3038-3051.
57. Beerli, P. and Felsenstein, J. (2001) Maximum likelihood estimation of a migration matrix and effective population sizes in n subpopulations by using a coalescent approach. *Proc. Natl. Acad. Sci. U.S.A.* 98, 4563-4568.
58. Whitlock, M.C. and McCauley, D.E. (1999) Indirect measures of gene flow and migration: F_{ST} not equal $1/(4Nm+1)$. *Heredity* 82, 117-125.
59. Saura, S. et al. (2014) Stepping stones are crucial for species' long-distance dispersal and range expansion through habitat networks. *J. Appl. Ecol.* 51, 171-182.

60. Fischer, M.C. et al. (2017) Estimating genomic diversity and population differentiation – an empirical comparison of microsatellite and SNP variation in *Arabidopsis halleri*. *BMC Genomics* 18, <https://doi.org/10.1186/s12864-016-3459-7>.
61. Hodel, R.G.J. et al. (2017) Adding loci improves phylogeographic resolution in red mangroves despite increased missing data: comparing microsatellites and RAD-Seq and investigating loci filtering. *Sci. Rep.* 7, <https://doi.org/10.1038/s41598-017-16810-7>.
62. Gillespie, R.G. et al. (2012) Long-distance dispersal: a framework for hypothesis testing. *Trends Ecol. Evol.* 27, 47-56.
63. Faillettaz, R. et al. (2018) Larval fish swimming behavior alters dispersal patterns from Marine Protected Areas in the North-Western Mediterranean Sea. *Front. Mar. Sci.* 5, <https://doi.org/10.3389/fmars.2018.00097>.
64. Nanninga, G.B. and Manica, A. (2018) Larval swimming capacities affect genetic differentiation and range size in demersal marine fishes. *Mar. Ecol. Prog. Ser.* 589, 1-12.
65. Carlton, J.T. et al. (2017) Tsunami-driven rafting: Transoceanic species dispersal and implications for marine biogeography. *Science* 357, 1402.
66. Shulzitski, K. et al. (2016) Encounter with mesoscale eddies enhances survival to settlement in larval coral reef fishes. *Proc. Natl. Acad. Sci. U.S.A.* 113, 6928-6933.
67. Mora, C. et al. (2001) Dispersal of juvenile and adult reef fishes associated with floating objects and their recruitment into Gorgona Island Reefs, Colombia. *Bull. Mar. Sci.* 68, 557-561.
68. Castro, J.J. et al. (2001) A general theory on fish aggregation to floating objects: An alternative to the meeting point hypothesis. *Rev. Fish Biol. Fish.* 11, 255-277.
69. Macfarlane, C.B.A. et al. (2013) Dispersal of marine benthic invertebrates through ice rafting. *Ecol.* 94, 250-256.

499 70. Waters, C.G. et al. (2013) A methodology for recruiting a giant clam, *Tridacna maxima*,
500 directly to natural substrata: A first step in reversing functional extinctions? *Biol.Cons.* 160,
501 19-24.

502 71. Clarke Murray, C. et al. (2011) Recreational boating: a large unregulated vector
503 transporting marine invasive species. *Divers. Distrib.* 17, 1161-1172.

504 72. Andrello, M. et al. (2013) Low Connectivity between Mediterranean Marine Protected
505 Areas: A biophysical modeling approach for the Dusky Grouper *Epinephelus marginatus*.
506 *PLoS ONE* 8, DOI: 10.1371/journal.pone.0068564.

507 73. Harrison, H.B. et al. (2012) Larval export from marine reserves and the recruitment
508 benefit for fish and fisheries. *Curr. Biol.* 22, 1023-1028.

509 74. Le Port, A. et al. (2017) Temperate marine protected area provides recruitment subsidies
510 to local fisheries. *Proc. Royal. Soc. Lond. B Biol. Sci.* 284, DOI: 10.1098/rspb.2017.1300.

511 75. Stuart-Smith, R.D. et al. (2015) Thermal biases and vulnerability to warming in the
512 world's marine fauna. *Nature* 528, 88-92.

513 76. Kremer, A. et al. (2012) Long-distance gene flow and adaptation of forest trees to rapid
514 climate change. *Ecol. Lett.* 15, 378-392.

515 77. Bolnick, D.I. and Nosil, P. (2007) Natural selection in populations subject to a migration
516 load. *Evol.* 61, 2229-2243.

517 78. Swan, K.D. et al. (2016) Managing marine biodiversity: The rising diversity and
518 prevalence of marine conservation translocations. *Conserv. Lett.* 9, 239-251.

519 79. Bellwood, D.R. et al. (2012) Human activity selectively impacts the ecosystem roles of
520 parrotfishes on coral reefs. *Proc. Natl. Acad. Sci. U.S.A* 279, DOI: 10.1098/rspb.2011.1906.

521 80. Pelc, R.A. et al. (2010) Detecting larval export from marine reserves. *Proc. Natl. Acad.*
522 *Sci. U.S.A.* 107, 18266-18271.

81. De Leo, G.A. and Micheli, F. (2015) The good, the bad and the ugly of marine reserves for fishery yields. *Philos. Trans. R. Soc. B Biol. Sci.* 370, DOI: 10.1098/rstb.2014.0276.
82. Claudet, J. et al. (2008) Marine reserves: size and age do matter. *Ecol. Lett.* 11, 481-489.
83. Barneche, D.R. et al. (2018) Fish reproductive-energy output increases disproportionately with body size. *Science* 360, 642-644.
84. Hopf, J.K. et al. (2016) Fishery consequences of marine reserves: short-term pain for longer-term gain. *Ecol. Appl.* 26, 818-829.
85. Halpern, B.S. et al. (2010) Placing marine protected areas onto the ecosystem-based management seascape. *Proc. Natl. Acad. Sci. U.S.A.* 107, 18312-18317.
86. Devillers, R. et al. (2014) Reinventing residual reserves in the sea: are we favouring ease of establishment over need for protection? *Aquat. Conserv.: Mar. Freshwat. Ecosyst.* 25, 480-504.
87. Cinner, J.E. et al. (2018) Gravity of human impacts mediates coral reef conservation gains. *Proc. Natl. Acad. Sci. U.S.A.* 115, DOI: 10.1073/pnas.1708001115.
88. Magris, R.A. et al. Biologically representative and well-connected marine reserves enhance biodiversity persistence in conservation planning. *Cons. Lett.* 11 <https://doi.org/10.1111/conl.12439>.
89. Treml, E.A. et al. (2015) Analyzing the (mis)fit between the institutional and ecological networks of the Indo-West Pacific. *Glob. Environ. Change* 31, 263-271.
90. Benestan, L. et al. (2015) RAD genotyping reveals fine-scale genetic structuring and provides powerful population assignment in a widely distributed marine species, the American lobster (*Homarus americanus*). *Mol. Ecol.* 24, 3299-3315.
91. Nielsen, E.E. et al. (2012) Gene-associated markers provide tools for tackling illegal fishing and false eco-certification. *Nat. Commun.* 3, DOI: 10.1038/ncomms1845.

- 547 92. Gaggiotti, O.E. (2017) Metapopulations of marine species with larval dispersal: A
548 counterpoint to Ilkka's Glanville fritillary metapopulations. *Ann. Zool. Fennici* 54, 97-112.
- 549 93. Bryan-Brown, D.N. et al. (2017) Patterns and trends in marine population connectivity
550 research. *Mar. Ecol. Prog. Ser.* 585, 243-256.
- 551 94. Kool, J.T. et al. (2013) Population connectivity: recent advances and new perspectives.
552 *Lands. Ecol.* 28, 165-185.
- 553 95. Calò, A. et al. (2013) A review of methods to assess connectivity and dispersal between
554 fish populations in the Mediterranean Sea. *Adv. Oceano. Limn.* 4, 150-175.
- 555 96. Gallego, A. et al. (2017) Bio-physical connectivity patterns of benthic marine species used
556 in the designation of Scottish nature conservation marine protected areas. *Ices J. Mar. Sci.* 74,
557 1797-1811.
- 558 97. Treml, E.A. et al. (2008) Modeling population connectivity by ocean currents, a graph-
559 theoretic approach for marine conservation. *Lands. Ecol.* 23, 19-36.
- 560 98. Treml, E.A. et al. (2015) Identifying the key biophysical drivers, connectivity outcomes,
561 and metapopulation consequences of larval dispersal in the sea. *Mov. Ecol.* 3, DOI:
562 10.1186/s40462-015-0045-6.
- 563 99. Planes, S. et al. (2009) Larval dispersal connects fish populations in a network of marine
564 protected areas. *Proc. Natl. Acad. Sci. U.S.A.* 106, 5693-5697.
- 565 100. Saenz-Agudelo, P. et al. (2011) Connectivity dominates larval replenishment in a coastal
566 reef fish metapopulation. *Proc. Royal. Soc. Lond. B Biol. Sci.* 278, 2954-2961.
- 567 101. Gandra, M. et al. (2018) Diel and seasonal changes in the spatial behaviour of a soft-
568 sediment fish (*Solea senegalensis*) inside a marine reserve. *Mar. Environ. Res.* 135, 82-92.
- 569 102. Starrs, D. et al. (2016) All in the ears: unlocking the early life history biology and spatial
570 ecology of fishes. *Biol. Rev.* 91, 86-105.

103. Calò, A. et al. (2013) A review of methods to assess connectivity and dispersal between fish populations in the Mediterranean Sea. *Adv. Oceano. Limn.* 4, 150 - 175.
104. Jones, G. (2015) Mission impossible: Unlocking the secrets of coral reef fish dispersal. In *Ecology of Fishes on Coral Reefs*. (Mora, C) Cambridge University Press, pp. 16-27. Cambridge, UK,.
105. Rousset, F. (2000) Genetic differentiation between individuals. *J. Evol. Biol.* 13, 58-62.
106. Leblois, R. et al. (2004) Influence of spatial and temporal heterogeneities on the estimation of demographic parameters in a continuous population using individual microsatellite data. *Genet.* 166, 1081-1092.
107. Gutenkunst, R.N. et al. (2009) Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. *PLoS Genet.* 5, <https://doi.org/10.1371/journal.pgen.1000695>.

Figure Legends

Figure 1: The spatial scale of sampling constraints dispersal estimates

(A) Boxplot representing the maximum dispersal distance and the sampling geographic extent across all studies. Central lines represent median values and whiskers first and third quartiles.

(B) Mean maximum dispersal distance increases with the maximal sampling geographic extent and (C) the pattern remains consistent among groups: fish, invertebrates and deep-sea organisms (>200 m). In (B) and (C), the color gradient displays the difference between maximal dispersal estimate and sampling geographic extent, with warmer colors (red) indicating that the maximum dispersal distance is closer to the maximal sampling geographic extent. The methods used to estimate connectivity include tracking, parentage analysis, assignment tests and isolation by distance (= demographic dispersal) described in Box 1 (see

B and C), biophysical models (potential dispersal), and genetic connectivity estimated from Fixation Index (F_{ST}). See supplementary method S1 and supplementary file S1 for details on the data used to generate the figure.

Figure 2: Potential long-distance dispersal processes and marine reserve benefits from

The main processes that contribute to long-dispersal distance are indicated with numbers and the main benefits due to long-dispersal distance are indicated with letters. The marine reserve is represented by a circle. (1) Active dispersal can drive larvae or adults far from the reserve boundaries, independently of the sea currents. (2) Larvae are pelagic and disperse passively due to currents. (3) They can associate with floating and drifting debris. (4) Translocation involves deliberately moving organisms from one site (“productive” reserves) to another (e.g. overexploited population). Thus, long distance dispersal can (A) increase biomass in fished areas far from the reserve, (B) potentially maintain species and genetic diversity across reserves, (C) maintain commercially and culturally important species that were the target of protection in the reserve.

Figure 3: Connectivity patterns in the global network of marine reserves.

(A) Map showing the neighbor distance for each marine reserve, i.e. the distance to the nearest marine reserves. To improve the visibility of the figure, we used both size and color of the circles to indicate the nearest-neighbor distance of each marine reserve. Small yellow circles indicate the most connected marine reserves (e.g. Scandinavian region or Australia) while large blue circles indicate the most isolated reserves (e.g. Western African coast). (B) Distribution of nearest-neighbor distances between marine reserves. The median and mean nearest-neighbor distances are 12 km and 65 km, respectively. The dashed red line indicates

620 the demographic median dispersal distance estimated for all organisms from the meta-analysis
621 (42 km). (C) The nearest-neighbor distance increases with the no-take surface area of marine
622 reserves (from 752 no-take marine reserves).

Box

Box 1: Methods for estimating long-distance dispersal in marine ecosystems

Methods to estimate marine dispersal [93-95] can be partitioned into three categories:

A. Potential dispersal (inferred from biophysical models)

Biophysical models can be used to simulate larval dispersal trajectories over large spatial and temporal scales (Table I) [96]. These models usually incorporate three elements: a physical model that simulates the ocean hydrodynamics, a particle tracking model that simulates the passive movement of virtual larvae, and optionally a coupled model that simulates the activity of the larvae when information on their ecology, behavior and physiology is available [97]. This third element is often lacking and it is therefore important to better understand the biology of marine larvae. Biophysical models are becoming increasingly complex and realistic, yet they always need to be validated with empirical data [98].

B. Realized dispersal (dispersal took place, but dispersers can or cannot successfully reproduce)

Specific dispersal events can be inferred using genetics. Parentage analyses identify dispersal events by using individual genotypes to assign juveniles to their parents [99]. This approach provides a snapshot of dispersal events over one generation. It requires considerable effort to sample and genotype a large number of juveniles and potential parents. Similarly, population genetic assignment tests use individual genotypes to assign individuals to their population of origin [48]. This approach relies on the occurrence of genetic structure among populations [100], but can also be applied in the absence of population genetic structure if populations are locally adapted [50]. Various tracking methods can also identify dispersal events. Acoustic telemetry can be used to observe the movement of individuals, often adults, providing the opportunity to directly observe dispersal [101]. Electronic and physical tags can provide

information on the movement of individuals. Some, however, have the drawback that individuals need to be recaptured to retrieve the data. Otolith analyses can also provide evidence of dispersal when the microchemistry or stable isotope composition of populations differ [102, 103]. When these approaches are applied to many individuals, it is possible to derive empirical distributions of dispersal kernel [104]. Yet the fact that individuals disperse does not necessary imply that they will successfully reproduce.

C. **Effective dispersal** (dispersal took place and dispersers successfully reproduced)

In the presence of **genetic isolation by distance** [105], it is possible to estimate dispersal at ecological timescales (tens of generations, [106]). This approach can be applied at the individual or population level. Another interesting avenue to infer dispersal is **cline analysis** [50]. Additional approaches based on **coalescent theory** [57] or the **site frequency spectrum** [107] go deeper back in time (tens to thousands of generations) and are therefore less relevant at ecological timescales.

Table I: Spatial extent and temporal resolution of the various methods used to estimate individual dispersal and connectivity in marine organisms.

Dispersal category	Method	Spatial extent			Temporal resolution	
		Low (1-40 km)	Medium (40-100 km)	Large (≥ 100 km)	Within generation	trans-generational
Potential	Biophysical		x	x	x	
Realized	Parentage	x	x	x		x
	Assignment	x	x	x	x	x
	Tracking	x	x	x	x	
	Otolith	x	x	x	x	x
Effective	Isolation by distance	x	x			x
	Cline analysis	x	x	x		x

Box 2

Translocation is the process by which living organisms are deliberately removed from one site for release in another. This definition excludes captive or cultivated organisms, sometimes genetically modified, that are massively released into the wild to support agriculture, fisheries, aquarium trade or pest control. The translocation process begins with the capture of wild organisms in a donor site and ends with post-release monitoring in the receiving site. Translocation has only recently become prominent in the oceans, particularly in coastal environments, where human impacts are the highest [67]. Translocations are equivalent to long-dispersal events.

Translocation in terrestrial environments is historically more common from non-protected to protected areas in order to prevent vulnerable organisms from being killed (e.g. African megafauna). In this scenario, protected areas are considered a sink, so the benefit is limited to individuals that are more likely to survive under protection. However, following the IUCN recommendations, conservation translocation must yield a measurable conservation benefit at the level of the population, species or ecosystem. Protected areas thus need to shift their role to become a source of translocated organisms and to provide long-distance benefits through human assistance. Marine reserves host more abundant and larger individuals, thus producing more larvae and juveniles than exploited areas [6, 19]. The challenge is now to capture these small larvae and juveniles, which are under high predation risk, to seed locally depleted or extinct populations elsewhere. This recently burgeoning strategy seems extremely promising. For instance, Palau, a small island nation created one of the largest marine reserves on the planet in 2015. Palau is now a sanctuary for marine animals that are globally endangered or

under severe threats like the Bumphead Parrotfish (*Bolbometopon muricatum*) which has critical and unique ecological functions in coral reef ecosystems [68]. The extremely high density of Bumphead Parrotfish in Palau induces massive spawning aggregations from which eggs can be collected and juveniles can be grown in tanks and then released at other sites where this species has been depleted (Figure I). Humans protect individuals through the period of high mortality and then release sub-adults in other reserves where populations have been exploited or are still being exploited to restore a certain density. This example highlights how marine reserves can play a pivotal role in long-distance translocations and broaden the geographic extent of their benefits in the near future.

Figure I. The translocation process for the threatened Bumphead Parrotfish (*Bolbometopon muricatum*) species from Palau. This island nation hosts the highest density of Bumphead Parrotfish worldwide (A) due to severe fishing restrictions. Massive spawning aggregations produce eggs (B) that can be caught without damage using nets (C). Larvae are then grown in optimal conditions to avoid mortality and juveniles (D) can be released to restore depleted or extinct local populations on overexploited reefs (E) but also revitalize a key functional role on coral reefs by bio-eroding dead corals (F). Photos from Tom Bowling (Biota Palau).

713 **Glossary**

714 **Admixture tracts:** Continuous blocks of the genome inherited from an admixed population.

715 **Benthic species:** Species that live and feed in or on the seabed.

716

717 **Blocks of identity by descent:** Continuous blocks of the genome that share the same alleles
718 inherited from a common ancestor.

719

720 **Cline analysis:** A framework that uses the relation between the genetic variation and the
721 geography or environment to estimate dispersal and selection.

722

723 **Coalescent theory:** A model that traces back gene variants from populations to their common
724 ancestor.

725

726 **Demersal species:** Species that live and feed near the bottom of the sea floor.

727

728 **Dispersal:** In this context, any movement of individuals or propagules from a source location
729 followed by successful immigration into a novel location with potential for gene flow.

730

731 **Demographic connectivity:** The process by which the dispersal of propagules, juveniles or
732 adults affects population growth and vital rates.

733

734 **Dispersal kernel:** Probability function describing the distribution of dispersal distances.

735

736 **Genetic connectivity:** A measure of gene flow and other evolutionary processes among
737 populations.

738

739 **Gene flow:** The exchange of genetic information among (sub)populations.

740

741 **Haplotype:** A combination of physically linked genetic variants on a single chromosome.

742

743 **Isolation by distance:** A pattern whereby genetic distance increases with geographic
744 distance. It can be used to estimate dispersal distance from population or individual genotype
745 data and regression analysis.

746

747 **Larval dispersal:** The dispersal of larvae from a spawning site to a settlement site.

748

749 **Next-generation sequencing:** Sequencing technologies that allows millions of DNA-
750 fragments to be sequenced in a single run.

751

752 **Pelagic larvae:** Larvae that spend time in the water column after hatching.

753

754 **Pelagic species:** Species living mainly in the water column.

755

756 **Sessile species:** Species that are fixed to a substratum for most of their life. Many sessile
757 species, however, have other stages in their life cycle, usually as eggs or larvae, that allow for
758 active or passive dispersal.

759

760 **Spillover:** The net movement of (adult and juvenile) organisms across the boundary of a
761 reserve into a fished area.

762

763 **Stepping-stone dispersal:** A dispersal process involving intermediate steps across several
764 generations.

765

766 **Single nucleotide polymorphism markers:** Molecular markers used to detect genetic
767 variation among individuals that correspond to a difference in a single DNA building block,
768 called a nucleotide.

769

770

771 Highlights and Outstanding questions inserted here for purposes of editorial markup.

772 **Highlights**

773 ▪ Marine dispersal estimates are limited by the spatial scale of sampling
774 design and therefore biased downwards;

775 ▪ Active larval behavior, oceanographic eddies and fronts, tsunamis,
776 marine debris and translocations are potentially important, but
777 overlooked, dispersal vectors over long distances;

778 ▪ The largest marine reserves have the highest potential for massive and
779 long-distance benefits, but are the most isolated ones;

780

- 781 ▪ Long-distance dispersal has important consequences for the design of
782 marine reserve networks;

783 Box 3: Outstanding questions Box

- 784 ▪ What proportion of the dispersal kernel of marine species are we
785 missing when we do not consider the long-distance (>40 km) dispersal?
- 786 ▪ Is the restricted spatial scale of sampling designs the only or main
787 cause of the limited geographical dispersal reported in the sea?
- 788 ▪ What are the quantitative effects of marine reserves at long distances
789 (>40 km)?
- 790 ▪ Is the unknown long-distance dispersal sufficient to connect large
791 isolated marine reserves?
- 792 ▪ To which extent future marine reserve networks should be composed of
793 few but large reserves instead of many small when we consider species
794 long-distance dispersal?

795

PROCESSES

1. Active dispersal
2. Passive dispersal by current
3. Drifting or floating debris
4. Translocation

BENEFITS

- A. Increase of biomass (Fisheries)
- B. Maintenance of biodiversity
- C. Maintenance of cultural species

○ Marine Reserve